

**Scheme 1** Synthesis of acceptor-substituted  $\beta$ -apo-8'-carotenoid compounds

**Table 1** Selected linear and nonlinear optical properties in chloroform of substituted  $\beta$ -apo-8'-carotenoids in Scheme 1

Compound	$\lambda_{\max}/\text{nm}$	$\epsilon/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$	$\mu/10^{-18} \text{ esu}$	$\beta/10^{-30} \text{ esu}$	$\beta(0)/10^{-30} \text{ esu}$	$\mu\text{-}\beta/10^{-48} \text{ esu}$
<b>1</b>	476	89 000	6.5	147	104	956
<b>2</b>	566	98 200	7.7	570	337	4389
<b>3</b>	502	95 700	7.0	220	148	1540

related compounds with chain lengths shorter than and equal to  $\beta$ -apo-8'-carotenal, **1** (Scheme 1), substituted with a series of donors and an aldehyde acceptor, and Ikeda and co-workers<sup>14</sup> have examined a series of retinal-derived compounds. In this contribution we report on the synthesis and second-order nonlinear optical properties of  $\beta$ -apo-8'-carotenal derivatives with terminal acceptor substituents, whose structures are more closely related to  $\beta$ -carotene than those in the previous studies.<sup>11–14</sup>

The new compounds were readily synthesized (Scheme 1) using standard Knoevenagel<sup>15</sup> conditions for the synthesis of **2** and Wadsworth–Emmons<sup>16</sup> conditions for **3**. The compounds were purified by chromatography on silica, recrystallized and then characterized by UV–VIS, <sup>1</sup>H and <sup>1</sup>H COSY NMR and mass spectroscopy, as well as elemental analysis.†  $\beta$  values were measured in chloroform solution by EFISH using 1.907  $\mu\text{m}$  fundamental radiation.<sup>6,7</sup>

Table 1 shows values of the optical absorption maximum ( $\lambda_{\max}$ ), the extinction coefficient ( $\epsilon$ ), the dipole moment ( $\mu$ ), first hyperpolarizability ( $\beta$ ), the zero frequency form of  $\beta$  [ $\beta(0)$ ] corrected for dispersion using a two-state model<sup>17</sup> and the dot product of  $\beta$  and  $\mu$  ( $\mu\text{-}\beta$ ), which of interest in poled polymer applications. Although these chromophores have no heteroatom donor, their hyperpolarizabilities are reasonably large, with each having a  $\beta(0)$  significantly larger than that of the prototypical stilbene compound 4-*N,N*-dimethylamino-4'-nitrostilbene (DANS) [ $\beta(0) = 55 \times 10^{-30} \text{ esu}$ ]. These rather large nonlinearities can be explained by the large intrinsic polarizability of the extended  $\pi$ -electron system. For polyenes, the HOMO–LUMO gap decreases with increasing conjugation length resulting from the energy of the LUMO being lowered and that of the HOMO being raised. In molecular orbital terms, a donor is a high lying occupied orbital, and thus, in long polyenes, the HOMO itself can act as a donor. As the length of the conjugated chain is increased, the HOMO becomes a more effective donor.

Dicyanovinyl groups are stronger acceptors than aldehydes, and this is reaffirmed by compound **2** having a  $\beta(0)$  value of over three times greater than for compound **1**. Nitro groups are also stronger acceptors than aldehydes, but on looking at the  $\beta(0)$  value for **3** it is not substantially greater than for **1**,

even although it is 6 atoms longer. This low value can be explained by the nitrophenyl acceptor containing a stabilizing aromatic group which hinders the formation of the charge-separated form of the molecule and hence reduces the polarizability and therefore  $\beta(0)$ .<sup>18</sup>

In summary,  $\beta(0)$  values of aldehyde **1**, dicyanovinyl **2** and nitrophenyl **3** acceptor-substituted  $\beta$ -apo-8'-carotenal chromophores measured by EFISH, are all large in view of the fact that no conventional donor is present, with **2** giving the largest value. The aromatic ring in compound **3** may hinder the formation of the charge-separated state, causing  $\beta(0)$  to be greatly reduced.

The work in this paper was performed, in part, by the Center for Space Microelectronics Technology, Jet Propulsion Laboratory, California Institute of Technology and was sponsored by the Strategic Defense Initiative Organization, Innovative Science and Technology Office and The Defense Advanced Research Projects Agency through grant no. 91-NC-146 administered by the Air Force Office of Scientific Research through agreements with the National Aeronautics and Space Administration (NASA). Support from the National Science Foundation (Grant CHE-9106689) is also gratefully acknowledged. S. G. thanks the National Research Council (NRC) and NASA for an NRC Resident Research Associateship at the Jet Propulsion Laboratory. We thank Brian Pierce and Joseph Perry for helpful discussions.

Received, 18th November 1992; Com. 2/06147C

## References

- S. R. Marder, D. N. Beratan and L.-T. Cheng, *Science*, 1991, **252**, 103.
- S. R. Marder, L.-T. Cheng and B. G. Tiemann, *J. Chem. Soc., Chem. Commun.*, 1992, 672.
- S. R. Marder, C. B. Gorman, L.-T. Cheng and B. G. Tiemann, *Proc. SPIE*, 1992, **1775**, in the press.
- J. J. P. Stewart, *J. Comput. Chem.*, 1989, **10**, 209.
- J. J. P. Stewart, *J. Comput. Chem.*, 1989, **10**, 221.
- B. F. Levine and C. G. Bethea, *Appl. Phys. Lett.*, 1974, **24**, 445.
- L.-T. Cheng, W. Tam, S. H. Stevenson, G. R. Meredith, G. Rikken and S. R. Marder, *J. Phys. Chem.*, 1991, **95**, 10631.
- E. Yang, A. C. Friedli, S. R. Marder and L.-T. Cheng, unpublished results.
- S. R. Marder, J. W. Perry, G. Bourhill, C. B. Gorman, B. G. Tiemann and K. Mansour, manuscript in preparation.
- J. B. VanBeek, F. Kajzar and A. C. Albrecht, *J. Chem. Phys.*, 1991, **95**, 6400.
- M. Blanchard-Desce, I. Ledoux, J.-M. Lehn, J. Malthête and J. Zyss, *J. Chem. Soc., Chem. Commun.*, 1988, 737.
- M. Barzoukas, M. Blanchard-Desce, D. Josse, J.-M. Lehn and J. Zyss, *Chem. Phys.*, 1989, **133**, 323.
- A. Slama-Schwok, M. Blanchard-Desce and J.-M. Lehn, *J. Chem. Phys.*, 1990, **94**, 3894.
- H. Ikeda, Y. Kawabe, T. Sakai, and K. Kawasaki, *Chem. Lett.*, 1989, **7**, 1285.
- B. S. Furniss, A. J. Hannaford, V. Rodgers, P. W. G. Smith and A. R. Tatchell, *Vogel's-Textbook of Practical Organic Chemistry*, Longman, New York, 4th edn., 1978, p. 490.
- W. S. Wadsworth and W. D. Emmons, *J. Am. Chem. Soc.*, 1961, **83**, 1733.
- J. L. Oudar and H. E. Le Person, *Opt. Commun.*, 1977, **66**, 2664.
- B. G. Tiemann, L.-T. Cheng and S. R. Marder, *J. Mater. Chem.*, to be submitted.

† Characterizing data for **2**: Yield, 15%. M.p., 184–186 °C. <sup>1</sup>H and <sup>1</sup>H COSY NMR (CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  1.03 (s, 6H), 1.47 (m, 2H), 1.62 (m, 2H), 1.72 (s, 3H), 1.99 (s, 3H), 2.00 (s, 3H), 2.02 (s, 3H), 2.03 (m, 2H), 2.27 (s, 3H), 6.15 (d, 1H, *J* 15.86 Hz), 6.18 (d, 1H, *J* 10.40 Hz), 6.24 (d, 1H, *J* 16.04 Hz), 6.31 (d, 1H, *J* 11.93 Hz), 6.39 (d, 1H, *J* 14.96 Hz), 6.54 (d, 1H, *J* 11.63 Hz), 6.64 (dd, 1H, *J* 14.76, 11.54 Hz), 6.68 (dd, 1H, *J* 13.91, 12.18 Hz), 6.78 (dd, 1H, *J* 14.93, 11.43 Hz), 6.86 (dd, 1H, *J* 13.75, 12.12 Hz), 6.80 (d, 1H, *J* 11.56 Hz), 6.81 (d, 1H, *J* 14.86 Hz), 7.26 (s, 1H). Satisfactory elemental analyses were obtained. High-resolution EIMS, *m/z* calc. for C<sub>33</sub>H<sub>40</sub>N<sub>2</sub>: 465.3270. Found: 465.3243.

Characterizing data for **3**: Yield, 37%. M.p., 296–300 °C. <sup>1</sup>H and <sup>1</sup>H COSY NMR (CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  1.03 (s, 6H), 1.47 (m, 2H), 1.62 (m, 2H), 1.72 (s, 3H), 1.98 (s, 3H), 1.99 (s, 3H), 2.00 (s, 3H), 2.03 (m, 2H), 2.07 (s, 3H), 6.14 (d, 1H, *J* 15.90 Hz), 6.16 (d, 1H, *J* 9.08 Hz), 6.21 (d, 1H, *J* 16.22 Hz), 6.28 (d, 1H, *J* 10.61 Hz), 6.35 (d, 1H, *J* 10.49 Hz), 6.37 (d, 1H, *J* 14.87 Hz), 6.47 (d, 1H, *J* 11.50 Hz), 6.52 (d, 1H, *J* 14.84 Hz), 6.63 (d, 1H, *J* 15.95 Hz), 6.66 (dd, 1H, *J* 10.74, 14.18 Hz), 6.70 (m, 1H), 6.71 (dd, 1H, *J* 11.26, 14.62 Hz), 6.72 (dd, 1H, *J* 10.80, 14.23 Hz), 7.09 (d, 1H, *J* 15.77 Hz), 7.56 (dm, 2H, *J* 8.61 Hz), 8.16 (dm, 2H, *J* 8.83 Hz). Satisfactory elemental analyses were obtained. High resolution EIMS, *m/z* calc. for C<sub>37</sub>H<sub>45</sub>NO<sub>2</sub>: 535.3442. Found: 535.3450.